

# Pre-eclampsia

Ben W J Mol, Claire T Roberts, Shakila Thangaratnam, Laura A Magee, Christianne J M de Groot, G Justus Hofmeyr



Pre-eclampsia affects 3–5% of pregnancies and is traditionally diagnosed by the combined presentation of high blood pressure and proteinuria. New definitions also include maternal organ dysfunction, such as renal insufficiency, liver involvement, neurological or haematological complications, uteroplacental dysfunction, or fetal growth restriction. When left untreated, pre-eclampsia can be lethal, and in low-resource settings, this disorder is one of the main causes of maternal and child mortality. In the absence of curative treatment, the management of pre-eclampsia involves stabilisation of the mother and fetus, followed by delivery at an optimal time. Although algorithms to predict pre-eclampsia are promising, they have yet to become validated. Simple preventive measures, such as low-dose aspirin, calcium, and diet and lifestyle interventions, show potential but small benefit. Because pre-eclampsia predisposes mothers to cardiovascular disease later in life, pregnancy is also a window for future health. A collaborative approach to discovery and assessment of the available treatments will hasten our understanding of pre-eclampsia and is an effort much needed by the women and babies affected by its complications.

## Introduction

Pre-eclampsia is a pregnancy-specific syndrome that affects 3–5% of pregnancies and is traditionally diagnosed when a pregnant woman presents with increased blood pressure and proteinuria.<sup>1</sup> Pre-eclampsia is one of the main causes of maternal, fetal, and neonatal mortality, especially in low-income and middle-income countries.<sup>2</sup> In this Seminar, we describe the current management of pre-eclampsia in terms of prediction, prevention, diagnosis, treatment, and long-term consequences. Our aim is to provide a guide for the optimal management of pre-eclampsia, both in low-resource and high-resource settings.

The acute clinical importance of pre-eclampsia lies in its relation to maternal and neonatal mortality and morbidity. When left untreated, pregnant women with pre-eclampsia have severe complications such as eclampsia, liver rupture, stroke, pulmonary oedema, or kidney failure, which can all be lethal.<sup>3</sup> Pre-eclampsia is also related to fetal growth restriction and preterm birth, either spontaneous or through iatrogenic delivery. Children born to mothers with pre-eclampsia have an increased risk of bronchopulmonary dysplasia and cerebral palsy, caused by preterm birth and being small for gestational age.<sup>4,5</sup> Pre-eclampsia decreases health-related quality of life and increases the risk of post-partum depression.<sup>6,7</sup>

The cause of pre-eclampsia is unclear. Some women are genetically predisposed to developing the disease which may run in families.<sup>8</sup> Robust associations have been identified between pre-eclampsia and gene variants involved in thrombophilia, inflammation, oxidative stress and the renin angiotensin system.<sup>9,10</sup> In a meta-analysis of studies to identify gene variants associated with pre-eclampsia, 22 variants were reproducible across studies with 7 remaining significant upon meta-analysis. However, thrombophilic gene variants in F2 and F5 have been consistently associated with the disease.<sup>11–13</sup> Interactions between maternal gene variants and genes encoding fetal HLA-C have been shown to predispose pregnancies to pre-eclampsia in white people, sub-Saharan

Africans, and the Chinese Han population, suggesting a role of an impaired immune tolerance in the pathogenesis of pre-eclampsia.<sup>14–16</sup> In women with pre-eclampsia, placental antiangiogenic factors are upregulated and disrupt the maternal endothelium, leading to an antiangiogenic state that can result in clinical signs of pre-eclampsia.<sup>17</sup>

## Definition of pre-eclampsia

The diagnostic criteria for pre-eclampsia were changed by the International Society for the Study of Hypertension in Pregnancy (ISSHP) in 2014.<sup>18</sup> ISSHP defines pre-eclampsia as de-novo hypertension present after 20 weeks of gestation combined with proteinuria (>300 mg/day), other maternal organ dysfunction, such as renal insufficiency, liver involvement, neurological or haematological complications, uteroplacental dysfunction, or fetal growth restriction. As proteinuria is no longer required in the new definition, proteinuric and non-proteinuric pre-eclampsia are now two separate categories.

Hypertension is defined as systolic blood pressure higher than 140 mm Hg or diastolic blood pressure higher than 90 mm Hg on two occasions that are 4–6 h apart.<sup>19–21</sup> Blood pressure should be measured in a seated and upright position or in a left lateral recumbent

Lancet 2016; 387: 999–1011

Published Online

September 3, 2015

[http://dx.doi.org/10.1016/S0140-6736\(15\)00070-7](http://dx.doi.org/10.1016/S0140-6736(15)00070-7)

S0140-6736(15)00070-7

The Robinson Research Institute, School of Paediatrics and Reproductive Health, University of Adelaide, SA, Australia (Prof B W J Mol PhD, Prof C T Roberts PhD); Women's Health Research Unit, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, London, UK (Prof S Thangaratnam PhD); BC Women's Hospital and Health Centre, Vancouver, BC, Canada (Prof L A Magee MD); Department of Obstetrics and Gynaecology, VU University Medical Center, Amsterdam, Netherlands

(Prof C J M de Groot PhD); and Effective Care Research Unit, University of the Witwatersrand, University of Fort Hare, and Eastern Cape Department of Health, East London, South Africa (Prof G J Hofmeyr DSc)

Correspondence to:

Prof B W J Mol, The Robinson Research Institute, School of Paediatrics and Reproductive Health, University of Adelaide, SA 5006, Australia  
[ben.mol@adelaide.edu.au](mailto:ben.mol@adelaide.edu.au)

## Search strategy and selection criteria

We searched PubMed and the Cochrane Library with the terms “pre-eclampsia” and “hypertension and pregnancy”, and cross-referenced them with the following terms: “epidemiology”, “definition”, “prediction”, “prevention”, “management”, “clinical trials”, “preconception care”, and “thrombophilia”. We restricted the search to studies done in humans and published in English. We limited our search to publications between January, 2010, and January, 2015, with a focus on publications after 2012. We also referred to older key publications. We then specifically looked at the themes of the review—ie, diagnostic studies, prognostic studies, intervention studies, and studies on long-term maternal risk.

position, using an appropriate size cuff and either manual or semi-automatic oscillometric devices that are validated for use in pre-eclampsia (Omron T9P or Omron MIT Elite devices).<sup>22,23</sup> Superimposed pre-eclampsia is diagnosed when women with underlying idiopathic hypertension present with one or more of the above features.

Women with proteinuria have high antenatal blood pressure, deliver at early stage of gestation, and often need operative delivery. Proteinuria is not an indicator of maternal morbidity or perinatal mortality, which could suggest a treatment paradox—ie, more aggressive treatment of women with severe pre-eclampsia might prevent complications. Optimum management (for example, of women with non-proteinuric pre-eclampsia or positive biomarkers) will be difficult to define without intervention studies that define the clinical effect of a diagnosis.<sup>24,25</sup>

### Prediction of pre-eclampsia

Although perfect prediction of pre-eclampsia has been a noble but hitherto elusive goal, distinction between women who are at low risk and high risk is possible. Strong risk factors are previous pre-eclampsia or hypertension in pregnancy, chronic kidney disease, hypertension, diabetes (type 1 or type 2), and autoimmune disorders, including systemic lupus erythematosus or antiphospholipid syndrome.<sup>20,26</sup> Moderate risk factors are first pregnancy, age 40 years or more, a pregnancy interval greater than 10 years, body-mass index of 35 kg/m<sup>2</sup> or more, polycystic ovarian syndrome, family history of pre-eclampsia, and multiple pregnancy.<sup>20,27</sup> Furthermore, women who have donated a kidney were twice as likely to have pre-eclampsia than matched women who had not donated a kidney.<sup>28</sup> However, in clinical practice, these factors predict just 30% of women who develop pre-eclampsia.<sup>29</sup>

Additional clinical and lifestyle factors that predict pre-eclampsia in early pregnancy include mean arterial pressure at 15 weeks of gestation, maternal birthweight, family history of coronary heart disease or pre-eclampsia, and vaginal bleeding for more than 5 days in the current pregnancy.<sup>30</sup> A previous single miscarriage with the same partner, time to conception of at least 12 months, and high fruit consumption were associated with a reduced risk of pre-eclampsia, but the combination of these factors resulted in only modest risk prediction (area under the receiver-operating characteristic curve [AUC] 0.71).<sup>30</sup>

Biomarkers in maternal blood have modest predictive potential in early pregnancy or have not been replicated across populations. Serum concentrations of maternal placental growth factor (proangiogenic) decreased in the blood 5 weeks before diagnosis of pre-eclampsia whereas those of soluble fms-like tyrosine kinase 1 (antiangiogenic) increased. However, the clinical value of these biomarkers has not been assessed in appropriately designed intervention studies.<sup>31–33</sup>

The combination of late first-trimester uterine artery Doppler ultrasound examination, placental growth factor, and pregnancy-associated plasma protein-A in maternal blood predict early-onset pre-eclampsia (sensitivity 93% [95% CI 76–98%]; specificity 95% [94–96%]), but this model still needs validation.<sup>34,35</sup> Results of a meta-analysis confirmed the accuracy of first-trimester uterine artery Doppler ultrasound for the prediction of early-onset pre-eclampsia (sensitivity 48%; specificity 92%).<sup>36</sup> The concentrations of circulating placental growth factor, vascular endothelial growth factor soluble fms-like tyrosine kinase 1, and soluble endoglin differ significantly before 30 weeks of gestation in women who developed pre-eclampsia, but the test accuracy of these markers is too poor to allow their use in clinical practice.<sup>37</sup>

In a large multicentre US study,<sup>38</sup> clinical risk factors combined with a change in maternal blood concentrations of angiogenic factors between first and early second trimester predicted early-onset pre-eclampsia (sensitivity 88%; specificity 80%). Results of another study showed that a combination of 11 biomarkers, clinical risk factors, and uterine artery Doppler ultrasound at 20 weeks of gestation had very good accuracy for prediction of pre-eclampsia (AUC 0.90 [95% CI 0.79–1.0]),<sup>39</sup> whereas investigators who took a metabolomics approach<sup>40</sup> found 14 metabolites at 15 weeks of gestation were predictive of pre-eclampsia (sensitivity 70%; specificity 95%). All these models need validation before they can be used in clinical practice.

### Prevention of pre-eclampsia

With reliable risk prediction for pre-eclampsia on the horizon, interventions to prevent pre-eclampsia become more important (table 1). Aspirin is the drug of choice for the prevention of pre-eclampsia, based on the findings of an individual patient data (IPD) meta-analysis that showed moderate benefit of aspirin (RR 0.90, 95% CI 0.84–0.97).<sup>41</sup> Other pharmacological drugs, such as heparin and dalteparin, show promising effects in women who are at increased risk for pre-eclampsia, but unlike trials of aspirin, the studies were too small to draw definite conclusions.<sup>42,51,52</sup>

Low dietary calcium and low serum calcium concentrations are associated with pre-eclampsia.<sup>53</sup> In women with low dietary calcium intake, high-dose calcium supplementation reduces pre-eclampsia (RR 0.36, 95% CI 0.20–0.65).<sup>43</sup> Although calcium supplementation is not recommended in women with normal dietary calcium intake, WHO recommends calcium supplementation (1.5–2 g daily) in the second half of pregnancy for women with low dietary calcium intake.<sup>54</sup> This recommendation applies to most settings in which the main dietary source of calcium is grains, such as wheat or maize, and is based on the systematic review of high-dose calcium supplementation during pregnancy.<sup>55</sup> The hypothesis that dietary calcium supplementation in early pregnancy might effectively

Population		Relative risk (95% CI)
<b>Pharmacological interventions</b>		
Aspirin	Women at risk of pre-eclampsia, gestational hypertension, or fetal growth restriction	0.90 (0.84–0.97) <sup>41</sup>
Low molecular weight heparin	Women at risk of placental dysfunction (eg, past history of pre-eclampsia, renal disease, placental abruption, fetal death, or fetal growth restriction)	0.47 (0.22–1.03) <sup>42</sup>
<b>Non-pharmacological interventions</b>		
Calcium	Women with low dietary calcium intake	0.36 (0.20–0.65) <sup>43</sup>
Vitamin C and E	Women at low, moderate, or high risk of pre-eclampsia	1.00 (0.92–1.09) <sup>44</sup>
Magnesium	Women at normal or high risk of pre-eclampsia	0.87 (0.58–1.3) <sup>45</sup>
L-arginine with antioxidants	Women at risk of pre-eclampsia	0.34 (0.21–0.55) <sup>46</sup>
Vitamin D and calcium	Women at any risk of pre-eclampsia	0.67 (0.33–1.4) <sup>47</sup>
Diet and lifestyle	Women at any risk of pre-eclampsia;* women who are overweight or obese; women who eat vegetables, fruits, and vegetable oils	0.74 (0.60–0.92); <sup>48</sup> 1.03 (0.71–1.5); <sup>49</sup> 0.72 (0.62–0.85) <sup>50</sup>

\*Includes women with mild gestational diabetes on insulin.

**Table 1: Interventions for prevention of pre-eclampsia**

prevent pre-eclampsia is being tested in a randomised trial in women with previous pre-eclampsia.<sup>56</sup>

Dietary supplementation with vitamin C and vitamin E (RR 1.00, 95% CI 0.92–1.09) or magnesium (0.87, 0.58–1.32) does not reduce the risk of pre-eclampsia.<sup>44,45</sup> Nutritional supplementation with marine oil or other long-chain polyunsaturated fatty acids is not effective in the prevention of pre-eclampsia either.<sup>57</sup> Vitamin D insufficiency is associated with an increased risk of gestational diabetes, pre-eclampsia, and small size for gestational age, but prophylactic vitamin D supplementation has only been assessed in one randomised controlled trial (0.67, 0.33–1.35).<sup>47,58</sup>

In a large randomised controlled trial of women at high risk of pre-eclampsia, the nitric oxide precursor L-arginine, reduced the risk of pre-eclampsia when given in combination with antioxidants (RR 0.17, 95% CI 0.12–0.21),<sup>46</sup> a finding that was later confirmed in a meta-analysis (0.34, CI 0.21–0.55).<sup>59</sup> Further studies of women at low risk are needed before supplementation of L-arginine in pregnancy can be routinely recommended.<sup>60</sup>

In response to the threat of the global obesity epidemic, the focus on healthy lifestyles to prevent pre-eclampsia is increasing. In a large cohort of nulliparous women, a diet rich in vegetables, fruits, and vegetable oils was associated with a reduced risk of pre-eclampsia (RR 0.72, 95% CI 0.62–0.85).<sup>50</sup> Systematic reviews<sup>48,61</sup> have shown that diet and lifestyle interventions can reduce the risk of pre-eclampsia in pregnant women, including women with gestational diabetes although this effect was not confirmed in a more recent randomised controlled trial<sup>49</sup> of diet and lifestyle interventions in pregnant women who are overweight or obese. In summary, treatment with aspirin is the only intervention to prevent pre-eclampsia for which robust evidence exists, but its effect is not large. Except for calcium supplementation in women with low dietary calcium intake, all other preventive interventions need

further assessment and should not be prescribed outside the context of clinical trials.

### Clinical presentation of pre-eclampsia

The clinical presentation of pre-eclampsia is varied.<sup>1</sup> Women are mostly asymptomatic, and the disease is often diagnosed during routine antenatal care. Maternal adverse outcomes are recorded in 10% of women with pre-eclampsia, whereas this risk increases to 15% in women with early-onset disease.<sup>62</sup>

The clinical presentation and findings could be indicative of the underlying multisystem morbidity. Women with severe pre-eclampsia might present with symptoms such as headache, visual disturbances (including blindness), epigastric pain, or nausea and vomiting. Neurological complications include eclamptic seizures, stroke, or reversible ischaemic neurological deficit, cortical blindness, retinal detachment, and posterior reversible encephalopathy. Hepatic involvement manifests as liver dysfunction, haematoma, or rupture, and renal involvement includes acute renal insufficiency requiring dialysis. Cardiorespiratory complications include myocardial ischaemia or infarction and pulmonary oedema. Women might also present with disseminated intravascular coagulation or placenta-related complications, such as abruption.

Severe pre-eclampsia could also be manifest as HELLP syndrome, characterised by microangiopathic haemolytic anaemia, hepatic dysfunction, and thrombocytopenia, with or without proteinuria or severe hypertension. HELLP syndrome often has an acute onset, with rapid deterioration of the maternal condition, and a third of cases present before 28 weeks of gestation.<sup>63</sup> Fetal complications include growth restriction, stillbirth, neonatal death, and prematurity-associated complications from early delivery.

Because of the heterogeneous clinical presentation of severe pre-eclampsia, many other disorders should be considered before definitive diagnosis (table 2).

	Presentation	Differential diagnosis
Central nervous system	Seizures, headache	Epilepsy, subarachnoid haemorrhage, hypoglycaemia, thrombotic thrombocytopenic purpura, hypertensive encephalopathy, central venous sinus thrombosis, local anaesthetic toxicity (epidural), amniotic fluid embolism, cerebral systemic lupus erythematosus, idiopathic intracranial hypertension
Renal	Proteinuria, hypertension, abnormal renal function tests, oliguria	Pyelonephritis, nephrotic syndrome, acute and chronic glomerulonephritis, lupus nephritis, haemolytic uraemic syndrome, interstitial nephritis
Vascular	Severe hypertension	Thyrotoxicosis, pheochromocytoma, Cushing's syndrome, white coat hypertension, hyperaldosteronism
Cardiorespiratory	Chest pain, dyspnoea, low oxygen saturation	Pulmonary oedema, pulmonary embolism, pneumonia, myocardial infarction or ischaemia, peripartum cardiomyopathy
Hepatic	Abnormal liver function tests, epigastric pain, nausea, vomiting	Acute fatty liver of pregnancy, viral hepatitis, drug-induced hepatotoxicity, acute pancreatitis, obstetric cholestasis, gastritis, hyperemesis gravidarum
Ophthalmological	Visual disturbances	Retinal detachment due to injury or eye diseases, retinal arterial or venous thrombosis due to vasculitis, trauma and other causes, retinal ischaemia, central serous retinopathy
Haematological	Bleeding, coagulation abnormality, disseminated intravascular coagulation, shock	Idiopathic thrombocytopenic purpura, thrombotic thrombocytopenic purpura, placental abruption, septic shock, acute fatty liver of pregnancy

**Table 2: Differential diagnosis of medical conditions with presentation similar to severe pre-eclampsia, by organ system involvement**

### Signs and symptoms

Clinicians routinely obtain information on the presence of symptoms associated with progression to severe disease. However, individual symptoms of pre-eclampsia such as headache (AUC 0.58, 95% CI 0.24–0.86), epigastric pain (0.70, 0.30–0.93), and visual disturbances (0.74, 0.33–0.94) do not adequately predict adverse maternal outcomes.<sup>64</sup> Chest pain and dyspnoea also have limited predictive value (0.64, 0.54–0.74) for composite adverse maternal outcomes.<sup>62</sup> Authors of a systematic review reported that mean arterial pressure of 140 mm Hg or higher or a blood pressure of 170/110 mm Hg or higher had limited accuracy for the prediction of eclampsia, placental abruption, and renal, neurological, and liver impairment (AUC for any adverse maternal outcome 0.68 [95% CI 0.29–0.92]).<sup>65</sup> Once pre-eclampsia has been diagnosed, blood pressure should be measured regularly in a day-assessment or inpatient setting, depending on the severity of the disorder.<sup>20,66</sup> Overall, history and physical examination to assess the severity of pre-eclampsia or predict complications have limited accuracy and should not be used alone to make management decisions such as delivery (table 3).

### Laboratory tests

The maternal spot urine estimate of the protein:creatinine ratio is promising for the detection of proteinuria in women with suspected pre-eclampsia. There is insufficient knowledge of how the protein:creatinine ratio should be used in clinical practice because test accuracy and prevalence across studies are heterogeneous.<sup>71</sup> The degree of proteinuria is not predictive for placental abruption or HELLP syndrome, whereas data on the capacity of proteinuria to predict eclampsia are conflicting.<sup>67,72</sup> Oxygen saturation predicts adverse maternal outcomes in the first 48 h after presentation fairly well (AUC 0.71, 95% CI 0.65–0.77).<sup>62</sup>

The monitoring of women with pre-eclampsia includes the assessment of haematological parameters (haemoglobin, platelets) and biochemical tests (hepatic and renal function) to track progression to severe disease and to diagnose deterioration of the disease. Individual tests, such as the detection of liver transaminases (AUC 0.79, 95% CI 0.51–0.93) perform fairly well, whereas others, such as the detection of a platelet count of less than  $100 \times 10^9$  per L (0.69, 0.63–0.75), serum creatinine (0.63, 0.57–0.69), and serum albumin (0.62, 0.56–0.68), have limited value in the prediction of complications.<sup>62,68</sup> Serum uric acid is a poor predictor of adverse outcomes and should not be measured.<sup>69</sup>

Routine assessment of the clotting profile is not necessary if the platelet count is more than  $100 \times 10^9$  platelets per L. Platelet count is not a sensitive indicator of coagulopathy. Outside pregnancy, neuraxial haematomas have not been reported with platelet counts of more than  $75 \times 10^9$  per L without platelet dysfunction or coagulopathy. Platelet transfusion (with or without other blood components) is indicated on the basis of platelet count, mode of delivery, presence of active bleeding, and coagulopathy.<sup>73</sup>

### Prediction models

Because all the above-mentioned tests have limited accuracy individually to predict complications, attempts have been made to integrate these tests into multivariable models. The full pre-eclampsia integrated estimate of risk (PIERS) multivariable model predicts composite adverse maternal outcome within 48 h in women admitted to hospital for pre-eclampsia (AUC ROC 0.88, 95% CI 0.84–0.92). In PIERS, the predictive factors are gestational age, platelet count, chest pain or dyspnoea, oxygen saturation, serum creatinine concentration, and aspartate transaminase concentration.<sup>62</sup> Surprisingly, blood pressure was not identified as a predictor of

	AUC (95% CI)	Sensitivity (95% CI)	Specificity (95% CI)	LR+ (95% CI)	LR- (95% CI)
<b>Symptoms</b>					
Headache					
Adverse maternal outcome*	0.58 (0.24–0.86) <sup>64</sup>	0.54 (0.27–0.79) <sup>64</sup>	0.59 (0.38–0.76) <sup>64</sup>	..	..
Epigastric pain					
Adverse maternal outcome*	0.70 (0.30–0.93) <sup>64</sup>	0.34 (0.22–0.50) <sup>65</sup>	0.83 (0.76–0.89) <sup>64</sup>	..	..
Visual disturbances					
Adverse maternal outcome*	0.74 (0.33–0.94) <sup>64</sup>	0.27 (0.07–0.65) <sup>65</sup>	0.81 (0.71–0.88) <sup>64</sup>	..	..
Nausea and vomiting					
Adverse maternal outcome*	0.54 (0.48–0.60) <sup>62</sup>	0.24 (0.21–0.27) <sup>60</sup>	0.87 (0.85–0.89) <sup>60</sup>	..	..
Chest pain or dyspnoea					
Adverse maternal outcome*	0.64 (0.54–0.74) <sup>62</sup>	..	..	..	..
<b>Examination</b>					
Blood pressure					
Adverse maternal outcome*	0.68 (0.29–0.92) <sup>64,65</sup>	..	..	..	..
Systolic blood pressure					
Adverse maternal outcome†	0.65 (0.59–0.70) <sup>62</sup>	..	..	..	..
Diastolic blood pressure					
Adverse maternal outcome*	0.63 (0.57–0.68) <sup>62</sup>	..	..	..	..
Mean arterial pressure					
Adverse maternal outcome*	0.65 (0.60–0.71) <sup>62</sup>	..	..	..	..
Oxygen saturation					
Adverse maternal outcome*	0.72 (0.67–0.78) <sup>62</sup>	..	..	..	..
<b>Investigation</b>					
24 h urine proteinuria increased by 2 g/24 h					
Eclampsia	..	..	..	0.41 (0.04–4.5) <sup>67</sup>	2.0 (0.83–4.6) <sup>67</sup>
Abrupton	..	..	..	1.1 (0.75–1.6) <sup>67</sup>	0.88 (0.42–1.9) <sup>67</sup>
HELLP syndrome	..	..	..	1.1 (0.74–1.6) <sup>67</sup>	0.86 (0.38–2.0) <sup>67</sup>
Dipstick proteinuria					
Adverse maternal outcome†	0.65 (0.59–0.71) <sup>62</sup>	..	..	..	..
Liver transaminases					
Adverse maternal outcome*	0.79 (0.51–0.93) <sup>68</sup>	..	..	..	..
Serum creatinine					
Adverse maternal outcome†	0.63 (0.57–0.69) <sup>62</sup>	..	..	..	..
Platelet count					
Adverse maternal outcome*	0.69 (0.63–0.75) <sup>62</sup>	..	..	..	..
Uric acid					
Eclampsia	..	..	..	2.1 (1.4–3.5) <sup>69</sup>	0.38 (0.18–0.81) <sup>69</sup>
Severe hypertension	..	..	..	1.7 (1.3–2.2) <sup>69</sup>	0.49 (0.38–0.64) <sup>69</sup>
Adverse maternal outcome†	0.59 (0.53–0.65) <sup>62</sup>	..	..	..	..
<b>Prediction models</b>					
Full PIERS model‡					
Adverse maternal outcome†	0.88 (0.84–0.92) <sup>62</sup>	..	..	..	..
Mini PIERS model§					
Adverse maternal outcome*	0.77 (0.74–0.80) <sup>70</sup>	..	..	..	..
<p>AUC=area under the curve. LR+=likelihood ratio of positive test. LR-=likelihood ratio of negative test. HELLP=haemolysis, elevated liver enzymes, low platelet count.  *Maternal adverse outcome to be present if any of the following occurs: maternal death, eclampsia, pulmonary oedema, abrupton, disseminated intravascular coagulation, renal failure, intracerebral haemorrhage, adult respiratory distress syndrome, and retinal detachment. †Maternal adverse outcome to be present if any of the following occurs: maternal mortality or one or more serious adverse events (central nervous system, cardiorespiratory, hepatic, renal, or haematological morbidity). ‡Based on gestational age, platelet count, symptoms such as chest pain or dyspnoea, oxygen saturation, and serum creatinine and aspartate transaminase concentrations. §Based on gestational age, headache or visual disturbances, chest pain or dyspnoea, vaginal bleeding with abdominal pain, systolic blood pressure, and dipstick proteinuria.</p>					
<b>Table 3: Accuracy of individual clinical tests and models in the prediction and diagnosis of maternal complications in women with pre-eclampsia</b>					

maternal complications, defined as the composite adverse maternal outcome in the model. The model findings might be affected by treatment paradox, where antihypertensive treatment or delivery might reduce high blood pressure-related events such as stroke, eclampsia, and transient ischaemic attack. Furthermore, the model might be affected by the low number of blood pressure-related components in the composite outcome. External validation of the PIERS model in an independent dataset showed high discrimination but poor calibration, perhaps because the external dataset was limited to women with severe pre-eclampsia.<sup>70</sup>

### Clinical management of women diagnosed with pre-eclampsia

Novel therapies for pre-eclampsia target various aspects of pre-eclampsia pathogenesis and are in development, yet the only cure for pre-eclampsia is delivery of the placenta (panel 1).<sup>86</sup> On the basis of consensus, severe hypertension or end-organ complications should be managed in an inpatient setting. Subspecialty consultation is advised to address underappreciation of the risks of pre-eclampsia. An early epidural catheter will attenuate pain-induced hypertension and enable neuraxial anaesthesia for emergency caesarean delivery, thereby avoiding difficult intubation or a hypertensive response to the intubation.<sup>74</sup> Bed rest does not prevent pre-eclampsia and is known to cause harm in general obstetrics.<sup>87</sup> To avoid potentially lethal pulmonary oedema, clinical practice has seen a trend towards fluid restriction, including the restrained use of fluids for oliguria, which has not been associated with an increase in renal failure. Avoidance of fluid preloading before neuraxial analgesia or anaesthesia is also recommended.<sup>88</sup>

International guidelines strongly recommend antihypertensive drugs for severe hypertension during pregnancy.<sup>20–22,74</sup> Repeated doses of nifedipine, intravenous hydralazine, or labetalol every 15–30 min all achieve blood pressure control in at least 80% of women.<sup>75–78</sup> Sufficiently powered randomised controlled trials are needed to compare regimens to achieve blood pressure control without overshoot. Because most severe pregnancy-related hypertension is not associated with end-organ dysfunction, lowering of blood pressure during a period of several hours is reasonable.<sup>79</sup> Contemporaneous use of nifedipine and magnesium sulphate is safe.<sup>89</sup>

Results of the recent CHIPS trial<sup>90</sup> showed that women with hypertension in pregnancy, be it chronic or pregnancy-induced, whose blood pressure was tightly controlled (target diastolic blood pressure 85 mm Hg) achieved a lower blood pressure (by 5 mm Hg) than women whose blood pressure was less tightly controlled (target diastolic blood pressure 100 mm Hg), resulting in similar rates of adverse perinatal outcome (adjusted odds ratio 0.98, 95% CI 0.74–1.3), a birthweight less than the tenth percentile (1.3, 0.93–1.8), and fewer women with severe maternal hypertension (0.56,

0.42–0.75) or serious maternal complications (2.0% vs 3.7%; 0.57, 0.26–1.27). Treatment of maternal hypertension benefits the mother, and although treatment might affect fetal growth, it does not increase illness or death of the infant.

Guidance for choice of the antihypertensive drug, including effects on fetal heart rate or neurodevelopment, is sparse.<sup>80,83</sup> Oral labetalol is not widely available in low-income and middle-income countries,<sup>81</sup> angiotensin converting enzyme inhibitors and receptor blockers are fetotoxic, prazosin can cause stillbirth, and atenolol might reduce fetal growth, the latter being controversial.<sup>74</sup> Individual variation in pre-eclampsia haemodynamics, assessed by cardiac output or peripheral vascular resistance, might interact with effects of antihypertensive drugs, but whether individualised therapy improves outcomes is unknown.

Intravenous magnesium sulphate is effective for treatment and prevention of eclampsia.<sup>84</sup> The high number of women with non-severe pre-eclampsia that need to be treated to prevent one seizure is an issue, especially in view of the cost of magnesium sulphate and its side-effects. Restricted use of magnesium sulphate or reduced magnesium sulphate dose or treatment duration, or both, have been suggested. The aim to give magnesium sulphate to women with severe (rather than any) pre-eclampsia might be associated with higher adverse outcome rates (ie, eclampsia, general anaesthesia, and neonatal morbidity).<sup>91</sup> Interest in magnesium sulphate dose reduction has been fuelled by fear of serious side-effects, which are not actually increased by magnesium sulphate (143 reports, 23916 women) and are infrequent (median <2%) in low-income and middle-income countries (24 studies, 9556 women).<sup>92,93</sup> In six randomised controlled trials (625 women), a reduction of magnesium sulphate dose or treatment duration, or both, resulted in outcomes similar to standard therapy, but sample sizes were too small for reliable conclusions.<sup>94</sup> Treatment of eclampsia with magnesium sulphate at the community level reduced recurrence (1 trial, 265 women),<sup>95</sup> and use of magnesium sulphate for pre-eclampsia is being tested in a cluster-randomised controlled trial (NCT01911494). Corticosteroids for HELLP help to improve laboratory parameters (11 trials, 550 women),<sup>85</sup> and the COHELLP trial (NCT00711841) will assess whether post-partum dexamethasone decreases maternal morbidity.<sup>96</sup>

All complications of pre-eclampsia can also occur post partum, particularly in the first 48 h. Intravenous labetalol and hydralazine are first-line drugs for the management of acute-onset, severe hypertension in the post-partum period, but oral nifedipine might also be considered as a first-line treatment post partum.<sup>97</sup>

### When to deliver the woman with pre-eclampsia

Because delivery of the placenta is the only cure for pre-eclampsia, optimal timing of delivery is crucial. The decision to deliver is based on the balance between the

**Panel 1: Management of women diagnosed with pre-eclampsia****Ante partum (irrespective of gestational age) and post partum (unless otherwise specified)***Place of care*<sup>74</sup>

- Inpatient care of patients with severe hypertension or maternal symptoms, signs, or abnormal laboratory tests
- Outpatient care can be considered, recognising that many women are not eligible and hospital readmission rates are high after home care

*Subspecialty consultation*<sup>74</sup>

- Obstetrics to ensure that pre-eclampsia risk is recognised and appropriate maternal and fetal surveillance is implemented
- Obstetric medicine (where available) to aid in management of hypertension and end-organ complications
- Anaesthesia to advise on monitoring of the woman and plan neuraxial analgesia or anaesthesia in labour to assist with blood pressure control and facilitate delivery by caesarean section (should it be necessary)

*Fluid management*<sup>74</sup>

- Restrict to a maximum of 80 mL/h when an intravenous drip is inserted

*Antihypertensive therapy*<sup>75-83</sup>

- For severe hypertension ( $\geq 160/110$  mm Hg), consider oral or parenteral agents that can be repeated in 30 min if blood pressure remains at  $\geq 160$  mm Hg systolic or  $\geq 110$  mm Hg diastolic: nifedipine capsule (10 mg orally without biting to a maximum of 30 mg); nifedipine tablet (10 mg orally to a maximum of 30 mg); hydralazine (5 mg intravenous bolus then 5–10 mg intravenous to a maximum of 45 mg if needed); labetalol (20 mg intravenous then, if needed, 40 mg then 80 mg to a maximum of 300 mg). Alternative oral drugs that can be repeated in 1 h (supported by less evidence in pregnancy): labetalol (200 mg orally); clonidine (0.1–0.2 mg orally);\*† captopril (only post partum 6.25–12.5 mg orally)\*

- For non-severe hypertension: methyldopa (500–2000 mg/dose in three or four divided doses); labetalol (300–2400 mg/dose in three or four divided doses); nifedipine (20–120 mg/dose once daily)

*Magnesium sulphate (MgSO<sub>4</sub>)*<sup>74,84</sup>

- Eclampsia treatment: 4 g intravenous (over 5 min), then 1 g/h intravenous; if patient is already receiving MgSO<sub>4</sub>, give additional 2–4 g intravenous (over 5 min) and increase infusion to 2 g/h intravenous
- Eclampsia prevention in women with pre-eclampsia: 4 g intravenous (over 5 min), then 1 g/h intravenous
- Fetal neuroprotection: 4 g intravenous (with or without 1 g/h until delivery or 24 h maximum) for women with imminent delivery at less than 34 weeks 0 days who do not otherwise qualify for eclampsia prevention or treatment

*Corticosteroids*<sup>85</sup>

- Antenatally only, to promote fetal pulmonary maturity when delivery is anticipated within the next 7 days and at less than 34 weeks 0 days
- HELLP syndrome (10 mg dexamethasone intravenous every 12 h for 48 h) if improvement in laboratory parameters alone will change management, such as eligibility for neuraxial anaesthesia or analgesia or platelet transfusion

*Platelet transfusion for HELLP syndrome*<sup>74</sup>

- Recommended for platelet counts  $< 20 \times 10^9$  per L,  $20 \times 10^9$ – $49 \times 10^9$  per L before caesarean, or  $\geq 50 \times 10^9$  per L (with or without packed erythrocytes) if patient is showing excessive active bleeding, platelet dysfunction, a rapidly diminishing platelet count, or coagulopathy

HELLP=haemolysis, elevated liver enzyme, low platelet. \*Captopril (25 mg) and clonidine (0.1 mg) are being compared in a post-partum randomised controlled trial (ClinicalTrials.gov, number NCT01761916) based on the effectiveness of these drugs in severe hypertension treatment outside pregnancy. †Clonidine therapy is not recommended during breastfeeding (<http://toxnet.nlm.nih.gov/newtoxnet/lactmed.htm>).

maternal and fetal risks of continuing the pregnancy and the neonatal risks of ending the pregnancy. In pre-eclampsia, the main driver is an assessment of the risk to the mother, but sometimes a growth-restricted child can become compromised and necessitate induction.

In a study that compared induction of labour and expectant management in women with pregnancy-induced hypertension or mild pre-eclampsia at term, induction of labour reduced the number of women with severe maternal hypertension, thus reducing the risk of severe maternal outcomes without affecting neonatal outcome; induction of labour also decreased the risk of caesarean section.<sup>98</sup> Women with an unripe cervix at term will, without induction, have a longer pregnancy and so will benefit most from an induction.<sup>99</sup> Oral misoprostol and mechanical induction with a transcervical balloon are safer than vaginal prostaglandins.<sup>100,101</sup>

For women with non-severe hypertensive disorders between 34 weeks and 37 weeks of gestation, immediate delivery, either through induction, or, if indicated, elective caesarean section, might reduce the already small risk of adverse maternal outcomes compared with expectant monitoring. However, immediate delivery increases the risk of neonatal respiratory distress syndrome. Therefore, routine immediate delivery does not seem justified, and a strategy of expectant monitoring until the clinical situation deteriorates can be considered until 37 weeks of gestation.<sup>102</sup> This strategy is only recommended if there are sufficient resources for safe monitoring while continuing the pregnancy.

Randomised studies that address risks at a gestational age of 34 weeks are scarce. In a South American multicentre trial,<sup>103</sup> women with early severe pre-eclampsia had an 8 day delay in birth after expectant management

compared with immediate delivery. Although neonatal outcome was comparable between the groups, expectant management was associated with an increase in placental abruption (RR 5.1, 95% CI 1.1–23). Timing of delivery in women with severe pre-eclampsia before 34 weeks remains a subject of research, but expectant management seems reasonable in well resourced settings. In women who present before 34 weeks of gestation with suspected pre-eclampsia, the ratio of circulating soluble fms-like tyrosine kinase 1 to placental growth factor predicts adverse outcomes occurring within 2 weeks.<sup>104</sup> The introduction of these biomarkers in clinical practice should be preceded by intervention studies.

In case of delivery before 34 weeks of gestation, infants benefit from a single course of antenatal corticosteroids to accelerate fetal lung maturation.<sup>105</sup> Exposure to antenatal corticosteroids after 34 weeks of gestation does not affect respiratory outcomes in infants with a subsequent late-preterm birth and is being assessed in the Antenatal Late Preterm Steroids (ALPS) trial (NCT01222247).<sup>106</sup> Babies born before 30 weeks of gestation are likely to benefit from the neuroprotective effect of antenatal treatment with magnesium sulphate (RR for cerebral palsy 0.68, 95% CI 0.54–0.87), although most women included in the underlying studies had premature rupture of membranes preceding spontaneous preterm birth.<sup>107</sup>

### Cardiovascular risk after pre-eclampsia

In 1964, Epstein showed that women with pre-eclampsia were at increased risk of developing cardiovascular disease later in life.<sup>108</sup> Until recently, women who have had a pregnancy complicated by hypertensive disorders, including pre-eclampsia, were not offered preventive measures. Evidence now overwhelmingly suggests that women who had pregnancy-related complications such as hypertensive disorders, gestational diabetes, intrauterine growth restriction, or preterm delivery have increased risk of cardiovascular disease later in life.<sup>107–112</sup> Pregnancy can therefore be considered as a window for future health.

Women with hypertension during pregnancy often have an unfavourable cardiovascular risk factor profile shortly after pregnancy. At 2 years after delivery, 30% of the women who had pre-eclampsia at term had hypertension and 25% had metabolic syndrome.<sup>113,114</sup> Pregnancy could possibly elicit metabolic syndrome, which then predisposes to vascular endothelial dysfunction.<sup>115</sup> The occurrence of gestational hypertension or gestational diabetes can re-emerge later in life as hypertension or type 2 diabetes.<sup>116</sup> Alternatively, pregnancy temporarily unmasks subclinical disease to return later in life. Women with a history of pre-eclampsia have an increased risk of microalbuminuria, with a prevalence similar to patients with type 1 diabetes.<sup>117</sup> Consequently, women who develop hypertension in pregnancy might qualify for a preventive strategy. This

group of women tend to also be motivated for lifestyle adjustment through smoking cessation and weight loss.<sup>118</sup> Several studies of post-partum caregivers revealed that there is limited knowledge of the association between hypertensive disorders in pregnancy and risk of cardiovascular disease, and only few health-care providers offer post-partum counselling for cardiovascular risk.<sup>119–121</sup>

Although there is an association between pre-eclampsia and cardiovascular disease in later life, it is unclear whether early screening and subsequent treatment reduce this increased cardiovascular risk. Enhancing awareness of cardiovascular risk has suggested to be effective by personal education postpartum.<sup>122</sup> Awareness is crucial but not sufficient for a healthy lifestyle post partum.<sup>123</sup> Recently, Berks and colleagues<sup>124</sup> reported that after pre-eclampsia, lifestyle interventions including exercise, dietary habits, and smoking cessation decreased cardiovascular risk by 4–13%. However, a history of gestational diabetes did not interact with the effect of a lifestyle intervention because rates of type 2 diabetes after 1 year were similar in women with and without gestational diabetes.<sup>125</sup> Before prevention measures for cardiovascular disease after pregnancy-related hypertension can be implemented, this knowledge gap must be filled.

In its 2011 guidelines for the prevention of cardiovascular disease in women, the American Heart Association<sup>126</sup> deemed a previous history of pre-eclampsia or gestational diabetes to be a major risk factor as part of its risk assessment system. The American Heart Association advises a yearly follow-up of blood pressure, lipid profile, and blood glucose concentration for women who had hypertension in pregnancy. However, intervention studies must be undertaken before such policies are implemented in clinical practice.

### Management of pre-eclampsia in low-resource settings

Pre-eclampsia and its serious consequences, including death, are many times more common in low-resource settings than in well resourced settings.<sup>20,127</sup> This epidemiological finding provides a model for seeking clues to the cause of pre-eclampsia and an obligation to direct global efforts to reduce mortality and morbidity from pre-eclampsia towards low-resource settings. Demographic risk factors for pre-eclampsia include young or advanced maternal age, family or personal history of hypertensive disorders, and poverty. Specifically in low-resource settings, an important strategy to reduce pre-eclampsia is the prevention of unintended pregnancy, particularly at the extremes of the reproductive age range, with high-quality, accessible family planning services.

Substantial progress is being made with smartphone-based systems to allow sophisticated prediction of pre-eclampsia risk in low-resource settings, using algorithms based on simple clinical data with or without



smartphone-based pulse oximetry.<sup>128</sup> In women who are admitted to hospital with pre-eclampsia, deterioration can now be predicted with the miniPIERS model, an adaptation of the PIERS model developed to predict the risk of adverse composite outcome in low-income and middle-income settings. The model integrates gestational age at the time of admission, headache or visual disturbances, chest pain or dyspnoea, vaginal bleeding with abdominal pain, systolic blood pressure, and dipstick proteinuria.<sup>74</sup>

Early detection and delivery before dangerous or irreversible complications arise is the mainstay of management of pre-eclampsia and is particularly difficult to implement in low-resource settings. Early pre-eclampsia is largely devoid of symptoms that would alert women to the need to seek medical care. Neither are there sufficiently sensitive predictors to allow identification of all women at risk. Regular routine antenatal assessments, including tests of blood pressure and proteinuria, are necessary for all pregnant women. Yet women with limited resources to invest in antenatal care are difficult to persuade to attend antenatal assessments when they feel well. Moreover, the principle of increased visit frequency in the last trimester when pre-eclampsia is most common has been abandoned in many low-resource settings in line with WHO's so-called basic antenatal care schedule of four antenatal visits for women at low risk of pre-eclampsia.<sup>129</sup> In view of evidence, including that from the original WHO Antenatal Care Trial,<sup>130</sup> that fewer visits are associated with increased perinatal mortality, we recommend that health services reassess the cost-effectiveness of more frequent antenatal visits in the third trimester (referred to as basic antenatal care plus).<sup>131</sup>

Dietary calcium intake is generally poor in low-resource settings.<sup>132</sup> The finding that pre-eclampsia was less common than expected in Mayan Indians living in Guatemala, who had relatively higher dietary calcium in their diets, led to the hypothesis that the links between poverty and pre-eclampsia might include dietary calcium deficiency.<sup>133</sup> Subsequent meta-analysis showed that calcium supplementation in the second half of pregnancy reduced pre-eclampsia in populations with low dietary calcium intake.<sup>46</sup>

The cost of 1.5–2 g calcium daily, as recommended by WHO, might be a limiting factor in low-resource settings.<sup>47</sup> A review of randomised trials assessing low-dose calcium supplementation (usually 500 mg daily) showed a consistent reduction in pre-eclampsia with calcium supplementation compared with placebo.<sup>134</sup> Pending review of the guidelines by WHO, 500 mg daily might be a reasonable alternative for women living in settings where diets are generally low in calcium if the recommended 1.5 g dosage is unachievable. The use of low-dose aspirin for women at high risk of pre-eclampsia is probably just as valid for women in low-resource settings as in well resourced settings.

#### Panel 2: Pre-eclampsia management strategies in low-resource settings

- Effective family planning services
- Algorithms for prediction of pre-eclampsia
- Calcium supplementation (1.5–2 g daily) for women with low dietary calcium intake
- Consider treatment with low-dose calcium (500 mg daily) if a high dose is unachievable
- Aspirin (75 mg daily) for women at high risk
- Frequent routine screening in the third trimester (basic antenatal care plus)
- Appropriate for prediction of disease progression
- Use of low-cost antihypertensive drugs, such as  $\alpha$ -methyl dopa
- Availability of magnesium sulphate for treatment of eclampsia
- Consideration of the cost vs benefit ratio of routine prophylaxis with magnesium sulphate in settings with limited resources for maternal monitoring
- Consider delivery in case of proteinuria and severe hypertension
- Use of a transcervical balloon with traction for labour induction to minimise uterine hyperstimulation,<sup>98</sup> with low-cost titrated oral misoprostol solution as second-line treatment<sup>99</sup>
- Conservative use of caesarean section
- Investment in health services

Once pre-eclampsia is diagnosed, the mainstay of treatment is to control blood pressure and deliver the placenta when the benefits of delivery outweigh the risks of conservative management. A substantial reduction in pre-eclampsia-related mortality could be achieved in low-income countries by widespread screening for hypertension and proteinuria and early delivery of women with severe disease.<sup>135</sup> Magnesium sulphate might reduce mortality but should not be the cornerstone of maternal mortality reduction programmes. Results of a cluster-randomised controlled trial<sup>136</sup> to assess increased use of antenatal corticosteroids in a low-resource setting showed that neonatal mortality did not decrease in infants with low birthweight, although neonatal mortality increased in the population overall.

Modelling of 2012 data has estimated that in sub-Saharan Africa, effective diagnosis, transfer, labour induction, and caesarean section would reduce annual maternal deaths from pre-eclampsia from 17 520 deaths with standard care at present to 4060 deaths, whereas further reduction with use of magnesium sulphate was limited from 4060 deaths to 3750 deaths.<sup>137</sup> Some modifications to reduce costs are possible, but there is no magic bullet. Reduction of maternal and perinatal mortality from pre-eclampsia in low-resource settings requires substantial investment in health systems, transport, health facilities, and skills training. Pragmatic

approaches specific to low-resource settings are shown in panel 2.

### Future research

Despite the scientific efforts (>17000 PubMed citations on pre-eclampsia since 2010) to increase knowledge of the aetiological and clinical aspects of pre-eclampsia, the incidence of the disease is not decreasing<sup>138</sup> and is still the main cause of maternal mortality, with significant adverse effects mainly in low-income and middle-income countries. Standardisation and collaboration in research will probably improve the use of resources. The Core Outcome Measures in Effectiveness Trials (COMET)<sup>139</sup> is developing a standardised outcome set for studies of hypertensive disorders in pregnancy. Such standardisation of study design in basic pre-eclampsia research has already been developed by the Global Pregnancy Collaboration (CoLab).<sup>140</sup> CoLab has also defined a pathway to optimise sample collection for placental research.<sup>141</sup> In another initiative, the Global Obstetric network (GONet) aims to coordinate and standardise the conduct of perinatal clinical trials<sup>142</sup> and the development of prediction models for pre-eclampsia through the global International Prediction of Pre-eclampsia IPD Collaborative Network, funded by the National Institute for Health Research. Adherence to standardised research protocols in a collaborative effort will result in larger datasets with standardised variables and will hasten the development of an understanding of the causes and prediction of pre-eclampsia and its complications. Development and assessment of targeted treatment strategies is an effort very much needed by the women and babies affected by pre-eclampsia.

### Contributors

BWJM structured the manuscript, did the search, wrote the introduction, and edited the manuscript. CTR wrote the section about the aetiology of pre-eclampsia and the section about prediction of pre-eclampsia. ST wrote the section about diagnosis and preventive treatment. LAM wrote the section on clinical management. CJMdG wrote the section on cardiovascular risk after pre-eclampsia. GJH wrote the section on management of pre-eclampsia in low-resource settings. All authors contributed to the manuscript and approved the final version.

### Declaration of interests

BWJM provides consultancy for ObsEva and has been an invited speaker on sponsored symposia and scientific meetings; all honoraria go to his institute. LAM was the principal investigator of the CHIPS trial and is a consultant to the PRE-EMPT project funded by the Bill & Melinda Gates Foundation. GJH is principal investigator for the Calcium and Pre-eclampsia Study funded by the University of British Columbia, a grantee of the Bill & Melinda Gates Foundation. CTR, ST, and CJMdG declare no competing interests.

### References

- Ananth CV, Keyes KM, Wapner RJ. Pre-eclampsia rates in the United States, 1980-2010: age-period-cohort analysis. *BMJ* 2013; **347**: f6564.
- Saleem S, McClure EM, Goudar SS, et al, and the Global Network Maternal Newborn Health Registry Study Investigators. A prospective study of maternal, fetal and neonatal deaths in low- and middle-income countries. *Bull World Health Organ* 2014; **92**: 605–12.
- Souza JP, Gülmezoglu AM, Vogel J, et al. Moving beyond essential interventions for reduction of maternal mortality (the WHO Multicountry Survey on Maternal and Newborn Health): a cross-sectional study. *Lancet* 2013; **381**: 1747–55.
- Hansen AR, Barnés CM, Folkman J, McElrath TF. Maternal preeclampsia predicts the development of bronchopulmonary dysplasia. *J Pediatr* 2010; **156**: 532–36.
- Strand KM, Heimstad R, Iversen AC, et al. Mediators of the association between pre-eclampsia and cerebral palsy: population based cohort study. *BMJ* 2013; **347**: f4089.
- Prick BW, Bijlenga D, Jansen AJ, et al. Determinants of health-related quality of life in the postpartum period after obstetric complications. *Eur J Obstet Gynecol Reprod Biol* 2015; **185**: 88–95.
- Blom EA, Jansen PW, Verhulst FC, et al. Perinatal complications increase the risk of postpartum depression. The Generation R Study. *BJOG* 2010; **117**: 1390–98.
- Williams PJ, Broughton Pipkin F. The genetics of pre-eclampsia and other hypertensive disorders of pregnancy. *Best Pract Res Clin Obstet Gynaecol* 2011; **25**: 405–17.
- Rana S, Karumanchi SA, Lindheimer MD. Angiogenic factors in diagnosis, management, and research in preeclampsia. *Hypertension* 2014; **63**: 198–202.
- Jebbink J, Wolters A, Fernando F, Afink G, van der Post J, Ris-Stalpers C. Molecular genetics of preeclampsia and HELLP syndrome—a review. *Biochim Biophys Acta* 2012; **1822**: 1960–69.
- Buurma AJ, Turner RJ, Driessen JH, et al. Genetic variants in pre-eclampsia: a meta-analysis. *Hum Reprod Update* 2013; **19**: 289–303.
- Staines-Urias E, Paez MC, Doyle P, et al. Genetic association studies in pre-eclampsia: systematic meta-analyses and field synopsis. *Int J Epidemiol* 2012; **41**: 1764–75.
- Fong FM, Sahemey MK, Hamed G, et al. Maternal genotype and severe preeclampsia: a HuGE review. *Am J Epidemiol* 2014; **180**: 335–45.
- Hiby SE, Walker JJ, O’Shaughnessy KM, et al. Combinations of maternal *KIR* and fetal *HLA-C* genes influence the risk of preeclampsia and reproductive success. *J Exp Med* 2004; **200**: 957–65.
- Nakimuli A, Chazara O, Hiby SE, et al. A *KIR B* centromeric region present in Africans but not Europeans protects pregnant women from pre-eclampsia. *Proc Natl Acad Sci USA* 2015; **112**: 845–50.
- Long W, Shi Z, Fan S, et al. Association of maternal *KIR* and fetal *HLA-C* genes with the risk of preeclampsia in the Chinese Han population. *Placenta* 2015; **36**: 433–37.
- Venkatesha S, Toporsian M, Lam C, et al. Soluble endoglin contributes to the pathogenesis of preeclampsia. *Nat Med* 2006; **12**: 642–49.
- Tranquilli AL, Dekker G, Magee L, et al. The classification, diagnosis and management of the hypertensive disorders of pregnancy: A revised statement from the ISSHP. *Pregnancy Hypertens* 2014; **4**: 97–104.
- Magee LA, Helewa M, Moutquin JM, von Dadelszen P, and the Hypertension Guideline Committee, and the Strategic Training Initiative in Research in the Reproductive Health Sciences (STRRHHS) Scholars. Diagnosis, evaluation, and management of the hypertensive disorders of pregnancy. *J Obstet Gynaecol Can* 2008; **30** (suppl): S1–48.
- National Institute for Health and Clinical Excellence (NICE). Hypertension in pregnancy. NICE clinical guideline 107. London: RCOG Press, 2011.
- American College of Obstetricians and Gynecologists, and the Task Force on Hypertension in Pregnancy. Hypertension in Pregnancy. *Obstet Gynecol* 2013; **122**: 1122–31.
- Brown MA, Roberts L, Davis G, Mangos G. Can we use the Omron T9P automated blood pressure monitor in pregnancy? *Hypertens Pregnancy* 2011; **30**: 188–93.
- Nouwen E, Sniijder M, van Montfrans G, Wolf H. Validation of the Omron M7 and Microlife 3BTO-A blood pressure measuring devices in preeclampsia. *Hypertens Pregnancy* 2012; **31**: 131–39.
- Glasziou P, Chalmers I, Rawlins M, McCulloch P. When are randomised trials unnecessary? Picking signal from noise. *BMJ* 2007; **334**: 349–51.
- Staff AC, Benton SJ, von Dadelszen P, et al. Redefining preeclampsia using placenta-derived biomarkers. *Hypertension* 2013; **61**: 932–42.
- Bramham K, Parnell B, Nelson-Piercy C, Seed PT, Poston L, Chappell LC. Chronic hypertension and pregnancy outcomes: systematic review and meta-analysis. *BMJ* 2014; **348**: g2301.
- Roos N, Kieler H, Sahlin L, Ekman-Ordeberg G, Falconer H, Stephansson O. Risk of adverse pregnancy outcomes in women with polycystic ovary syndrome: population based cohort study. *BMJ* 2011; **343**: d6309.

- 28 Garg AX, Nevis IF, McArthur E, et al, and the DONOR Network. Gestational hypertension and preeclampsia in living kidney donors. *N Engl J Med* 2015; **372**: 124–33.
- 29 Leslie K, Thilaganathan B, Papageorghiou A. Early prediction and prevention of pre-eclampsia. *Best Pract Res Clin Obstet Gynaecol* 2011; **25**: 343–54.
- 30 North RA, McCowan LME, Dekker GA, et al. Clinical risk prediction for pre-eclampsia in nulliparous women: development of model in international prospective cohort. *BMJ* 2011; **342**: d1875.
- 31 Levine RJ, Maynard SE, Qian C, et al. Circulating angiogenic factors and the risk of preeclampsia. *N Engl J Med* 2004; **350**: 672–83.
- 32 Hagmann H, Thadhani R, Benzing T, Karumanchi SA, Stepan H. The promise of angiogenic markers for the early diagnosis and prediction of preeclampsia. *Clin Chem* 2012; **58**: 837–45.
- 33 Chappell LC, Duckworth S, Seed PT, et al. Diagnostic accuracy of placental growth factor in women with suspected preeclampsia: a prospective multicenter study. *Circulation* 2013; **128**: 2121–31.
- 34 Allen RE, Rogozinska E, Cleverly K, Aquilina J, Thangaratinam S. Abnormal blood biomarkers in early pregnancy are associated with preeclampsia: a meta-analysis. *Eur J Obstet Gynecol Reprod Biol* 2014; **182**: 194–201.
- 35 Poon LCY, Kametas NA, Maiz N, Akolekar R, Nicolaides KH. First-trimester prediction of hypertensive disorders in pregnancy. *Hypertension* 2009; **53**: 812–18.
- 36 Velauthar L, Plana MN, Kalidindi M, et al. First-trimester uterine artery Doppler and adverse pregnancy outcome: a meta-analysis involving 55,974 women. *Ultrasound Obstet Gynecol* 2014; **43**: 500–07.
- 37 Kleinrouweler CE, Wiegerinck MM, Ris-Stalpers C, et al, and the EBM CONNECT Collaboration. Accuracy of circulating placental growth factor, vascular endothelial growth factor, soluble fms-like tyrosine kinase 1 and soluble endoglin in the prediction of pre-eclampsia: a systematic review and meta-analysis. *BJOG* 2012; **119**: 778–87.
- 38 Myatt L, Clifton RG, Roberts JM, et al, and the Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. Can changes in angiogenic biomarkers between the first and second trimesters of pregnancy predict development of pre-eclampsia in a low-risk nulliparous patient population? *BJOG* 2013; **120**: 1183–91.
- 39 Kenny LC, Black MA, Poston L, et al. Early pregnancy prediction of preeclampsia in nulliparous women, combining clinical risk and biomarkers: the Screening for Pregnancy Endpoints (SCOPE) international cohort study. *Hypertension* 2014; **64**: 644–52.
- 40 Kenny LC, Broadhurst DI, Dunn W, et al, and the Screening for Pregnancy Endpoints Consortium. Robust early pregnancy prediction of later preeclampsia using metabolomic biomarkers. *Hypertension* 2010; **56**: 741–49.
- 41 Askie LM, Duley L, Henderson-Smart DJ, Stewart LA, and the PARIS Collaborative Group. Antiplatelet agents for prevention of pre-eclampsia: a meta-analysis of individual patient data. *Lancet* 2007; **369**: 1791–98.
- 42 Dodd JM, McLeod A, Windrim RC, Kingdom J. Antithrombotic therapy for improving maternal or infant health outcomes in women considered at risk of placental dysfunction. *Cochrane Database Syst Rev* 2013; **7**: CD006780.
- 43 Hofmeyr GJ, Lawrie TA, Atallah AN, Duley L, Torloni MR. Calcium supplementation during pregnancy for preventing hypertensive disorders and related problems. *Cochrane Database Syst Rev* 2014; **6**: CD001059.
- 44 Conde-Agudelo A, Romero R, Kusanovic JP, Hassan SS. Supplementation with vitamins C and E during pregnancy for the prevention of preeclampsia and other adverse maternal and perinatal outcomes: a systematic review and meta-analysis. *Am J Obstet Gynecol* 2011; **204**: 503. e1–12.
- 45 Makrides M, Crosby DD, Bain E, Crowther CA. Magnesium supplementation in pregnancy. *Cochrane Database Syst Rev* 2014; **4**: CD000937.
- 46 Dorniak-Wall T, Grivell RM, Dekker GA, Hague W, Dodd JM. The role of L-arginine in the prevention and treatment of pre-eclampsia: a systematic review of randomised trials. *J Hum Hypertens* 2014; **28**: 230–35.
- 47 De-Regil LM, Palacios C, Ansary A, Kulier R, Peña-Rosas JP. Vitamin D supplementation for women during pregnancy. *Cochrane Database Syst Rev* 2012; **2**: CD008873.
- 48 Thangaratinam S, Rogozinska E, Jolly K, et al. Effects of interventions in pregnancy on maternal weight and obstetric outcomes: meta-analysis of randomised evidence. *BMJ* 2012; **344**: e2088.
- 49 Dodd JM, Turnbull D, McPhee AJ, et al, and the LIMIT Randomised Trial Group. Antenatal lifestyle advice for women who are overweight or obese: LIMIT randomised trial. *BMJ* 2014; **348**: g1285.
- 50 Brantsaeter AL, Haugen M, Samuelsen SO, et al. A dietary pattern characterized by high intake of vegetables, fruits, and vegetable oils is associated with reduced risk of preeclampsia in nulliparous pregnant Norwegian women. *J Nutr* 2009; **139**: 1162–68.
- 51 Rodger MA, Hague WM, Kingdom J, et al, and the TIPPS Investigators. Antepartum dalteparin versus no antepartum dalteparin for the prevention of pregnancy complications in pregnant women with thrombophilia (TIPPS): a multinational open-label randomised trial. *Lancet* 2014; **384**: 1673–83.
- 52 de Vries JIP, van Pampus MG, Hague WM, Bezemer PD, Joosten JH, and the FRUIT Investigators. Low-molecular-weight heparin added to aspirin in the prevention of recurrent early-onset pre-eclampsia in women with inheritable thrombophilia: the FRUIT-RCT. *J Thromb Haemost* 2012; **10**: 64–72.
- 53 Ephraim RK, Osakunor DN, Denkyira SW, Eshun H, Amoah S, Anto EO. Serum calcium and magnesium levels in women presenting with pre-eclampsia and pregnancy-induced hypertension: a case-control study in the Cape Coast metropolis, Ghana. *BMC Pregnancy Childbirth* 2014; **14**: 390.
- 54 Hofmeyr GJ, Duley L, Atallah A. Dietary calcium supplementation for prevention of pre-eclampsia and related problems: a systematic review and commentary. *BJOG* 2007; **114**: 933–43.
- 55 Hofmeyr GJ, Lawrie TA, Atallah AN, Duley L. Calcium supplementation during pregnancy for preventing hypertensive disorders and related problems. *Cochrane Database Syst Rev* 2010; **8**: CD001059.
- 56 Hofmeyr GJ, Novikova N, Singata M, et al. Protocol 11PRT/4028: Long term calcium supplementation in women at high risk of pre-eclampsia: a randomised, placebo-controlled trial (PACTR201105000267371). <http://www.thelancet.com/protocol-reviews/11PRT-4028> (accessed April 7, 2015).
- 57 Makrides M, Duley L, Olsen SF. Marine oil, and other prostaglandin precursor, supplementation for pregnancy uncomplicated by pre-eclampsia or intrauterine growth restriction. *Cochrane Database Syst Rev* 2006; **3**: CD003402.
- 58 Aghajafari F, Nagulesapillai T, Ronksley PE, Tough SC, O'Beirne M, Rabi DM. Association between maternal serum 25-hydroxyvitamin D level and pregnancy and neonatal outcomes: systematic review and meta-analysis of observational studies. *BMJ* 2013; **346**: f1169.
- 59 Vadillo-Ortega F, Perichart-Perera O, Espino S, et al. Effect of supplementation during pregnancy with L-arginine and antioxidant vitamins in medical food on pre-eclampsia in high risk population: randomised controlled trial. *BMJ* 2011; **342**: d2901.
- 60 Meher S, Duley L. Nitric oxide for preventing pre-eclampsia and its complications. *Cochrane Database Syst Rev* 2007; **2**: CD006490.
- 61 Allen R, Rogozinska E, Sivarajasingam P, Khan KS, Thangaratinam S. Effect of diet- and lifestyle-based metabolic risk-modifying interventions on preeclampsia: a meta-analysis. *Acta Obstet Gynecol Scand* 2014; **93**: 973–85.
- 62 von Dadelszen P, Payne B, Li J, et al, and the PIERS Study Group. Prediction of adverse maternal outcomes in pre-eclampsia: development and validation of the fullPIERS model. *Lancet* 2011; **377**: 219–27.
- 63 Habli M, Eftekhari N, Wiebracht E, et al. Long-term maternal and subsequent pregnancy outcomes 5 years after hemolysis, elevated liver enzymes, and low platelets (HELLP) syndrome. *Am J Obstet Gynecol* 2009; **201**: 385. e1.
- 64 Thangaratinam S, Gallos ID, Meah N, Usman S, Ismail KM, Khan KS, and the TIPPS (Tests in Prediction of Pre-eclampsia's Severity) Review Group. How accurate are maternal symptoms in predicting impending complications in women with preeclampsia? A systematic review and meta-analysis. *Acta Obstet Gynecol Scand* 2011; **90**: 564–73.
- 65 Thangaratinam S, Datta A, Ismail KMK, Khan KSK. What is the accuracy of blood pressure in predicting complications in pre-eclampsia? *Arch Dis Child Fetal Neonatal Ed* 2011; **96**: 101.
- 66 Turnbull DA, Wilkinson C, Gerard K, et al. Clinical, psychosocial, and economic effects of antenatal day care for three medical complications of pregnancy: a randomised controlled trial of 395 women. *Lancet* 2004; **363**: 1104–09.

- 67 Thangaratinam S, Coomarasamy A, O'Mahony F, et al. Estimation of proteinuria as a predictor of complications of pre-eclampsia: a systematic review. *BMC Med* 2009; 7: 10.
- 68 Thangaratinam S, Koopmans CM, Iyengar S, et al, and the TIPPS (Tests in Prediction of Preeclampsia's Severity) Review Group. Accuracy of liver function tests for predicting adverse maternal and fetal outcomes in women with preeclampsia: a systematic review. *Acta Obstet Gynecol Scand* 2011; 90: 574–85.
- 69 Thangaratinam S, Ismail KM, Sharp S, Coomarasamy A, Khan KS, and the Tests in Prediction of Pre-eclampsia Severity review group. Accuracy of serum uric acid in predicting complications of pre-eclampsia: a systematic review. *BJOG* 2006; 113: 369–78.
- 70 Payne BA, Hutcheon JA, Ansermino JM, et al, and the miniPIERS Study Working Group. A risk prediction model for the assessment and triage of women with hypertensive disorders of pregnancy in low-resourced settings: the miniPIERS (Pre-eclampsia Integrated Estimate of RiSk) multi-country prospective cohort study. *PLoS Med* 2014; 11: e1001589.
- 71 Morris RK, Riley RD, Doug M, Deeks JJ, Kilby MD. Diagnostic accuracy of spot urinary protein and albumin to creatinine ratios for detection of significant proteinuria or adverse pregnancy outcome in patients with suspected pre-eclampsia: systematic review and meta-analysis. *BMJ* 2012; 345: e4342. DOI:10.1136/bmj.e4342.
- 72 Koopmans CM, Zwart JJ, Groen H, et al. Risk indicators for eclampsia in gestational hypertension or mild preeclampsia at term. *Hypertens Pregnancy* 2011; 30: 433–46.
- 73 Laskin S, Payne B, Hutcheon JA, et al. The role of platelet counts in the assessment of inpatient women with preeclampsia. *J Obstet Gynaecol Can* 2011; 33: 900–08.
- 74 Magee LA, Pels A, Helewa M, Rey E, von Dadelszen P, and the Canadian Hypertensive Disorders of Pregnancy (HDP) Working Group. Diagnosis, evaluation, and management of the hypertensive disorders of pregnancy. *Pregnancy Hypertens* 2014; 4: 105–45.
- 75 Duley L, Meher S, Jones L. Drugs for treatment of very high blood pressure during pregnancy. *Cochrane Database Syst Rev* 2013; 7: CD001449.
- 76 Lakshmi BS, Dasar P. Oral nifedipine versus intravenous labetalol in hypertensive urgencies and emergencies of pregnancy: a randomized clinical trial. *Obstet Med* 2012; 5: 171–75.
- 77 Saudan P, Billieux M-H, Pecheur A, Irion O, Savoldelli G, Boulvain M. OS014. Which first-line drug to control severe hypertension in pregnancy? A pilot study. *Pregnancy Hypertens* 2012; 2: 182.
- 78 Vermillion ST, Scardo JA, Newman RB, Chauhan SP. A randomized, double-blind trial of oral nifedipine and intravenous labetalol in hypertensive emergencies of pregnancy. *Am J Obstet Gynecol* 1999; 181: 858–61.
- 79 Firoz T, Magee LA, MacDonell K, et al, and the Community Level Interventions for Pre-eclampsia (CLIP) Working Group. Oral antihypertensive therapy for severe hypertension in pregnancy and postpartum: a systematic review. *BJOG* 2014; 121: 1210–18, discussion 1220.
- 80 Abalos E, Duley L, Steyn DW. Antihypertensive drug therapy for mild to moderate hypertension during pregnancy. *Cochrane Database Syst Rev* 2014; 2: CD002252.
- 81 Lalani S, Firoz T, Magee LA, et al. Pharmacotherapy for preeclampsia in low and middle income countries: an analysis of essential medicines lists. *J Obstet Gynaecol Can* 2013; 35: 215–23.
- 82 Orbach H, Matok I, Gorodischer R, et al. Hypertension and antihypertensive drugs in pregnancy and perinatal outcomes. *Am J Obstet Gynecol* 2013; 208: e1–6.
- 83 Gillon TE, Pels A, von Dadelszen P, MacDonell K, Magee LA. Hypertensive disorders of pregnancy: a systematic review of international clinical practice guidelines. *PLoS One* 2014; 9: e113715.
- 84 Altman D, Carroli G, Duley L, et al, and the Magpie Trial Collaboration Group. Do women with pre-eclampsia, and their babies, benefit from magnesium sulphate? The Magpie Trial: a randomised placebo-controlled trial. *Lancet* 2002; 359: 1877–90.
- 85 Woudstra DM, Chandra S, Hofmeyr GJ, Dowswell T. Corticosteroids for HELLP (hemolysis, elevated liver enzymes, low platelets) syndrome in pregnancy. *Cochrane Database Syst Rev*. 2010; 9: CD008148.
- 86 Everett TR, Wilkinson IB, Lees CC. Drug development in preeclampsia: a 'no go' area? *J Matern Fetal Neonatal Med* 2012; 25: 50–52.
- 87 Lorenz RP. What is new in bed rest in pregnancy? Best articles from the past year. *Obstet Gynecol* 2014; 124: 377–78.
- 88 WHO. WHO recommendations for prevention and treatment of pre-eclampsia and eclampsia. Geneva: World Health Organization, 2011.
- 89 Magee LA, Miremadi S, Li J, et al. Therapy with both magnesium sulfate and nifedipine does not increase the risk of serious magnesium-related maternal side effects in women with preeclampsia. *Am J Obstet Gynecol* 2005; 193: 153–63.
- 90 Magee LA, von Dadelszen P, Rey E, et al, and the CHIPS Study Group. Less-tight versus tight control of hypertension in pregnancy. *N Engl J Med* 2015; 372: 407–17.
- 91 Alexander JM, McIntire DD, Leveno KJ, Cunningham FG. Selective magnesium sulfate prophylaxis for the prevention of eclampsia in women with gestational hypertension. *Obstet Gynecol* 2006; 108: 826–32.
- 92 Bain ES, Middleton PF, Crowther CA. Maternal adverse effects of different antenatal magnesium sulphate regimens for improving maternal and infant outcomes: a systematic review. *BMC Pregnancy Childbirth* 2013; 13: 195.
- 93 Smith JM, Lowe RF, Fullerton J, Currie SM, Harris L, Felker-Kantor E. An integrative review of the side effects related to the use of magnesium sulfate for pre-eclampsia and eclampsia management. *BMC Pregnancy Childbirth* 2013; 13: 34.
- 94 Duley L, Matar HE, Almerie MQ, Hall DR. Alternative magnesium sulphate regimens for women with pre-eclampsia and eclampsia. *Cochrane Database Syst Rev* 2010; 8: CD007388.
- 95 PRE-EMPT. Community-level interventions for pre-eclampsia. <http://pre-empt.cfri.ca/treatment/clip-trial> (accessed April 7, 2015).
- 96 Katz L, Amorim M, Souza JP, Haddad SM, Cecatti JG, and the COHELLP Study Group. COHELLP: collaborative randomized controlled trial on corticosteroids in HELLP syndrome. *Reprod Health* 2013; 10: 28.
- 97 Committee on Obstetric Practice. Committee Opinion No. 623: Emergent therapy for acute-onset, severe hypertension during pregnancy and the postpartum period. *Obstet Gynecol* 2015; 125: 521–25.
- 98 Koopmans CM, Bijlenga D, Groen H, et al, and the HYPITAT study group. Induction of labour versus expectant monitoring for gestational hypertension or mild pre-eclampsia after 36 weeks' gestation (HYPITAT): a multicentre, open-label randomised controlled trial. *Lancet* 2009; 374: 979–88.
- 99 Tajik P, van der Tuuk K, Koopmans CM, et al. Should cervical favourability play a role in the decision for labour induction in gestational hypertension or mild pre-eclampsia at term? An exploratory analysis of the HYPITAT trial. *BJOG* 2012; 119: 1123–30.
- 100 Jozwiak M, Oude Rengerink K, Benthem M, et al, and the PROBAAT Study Group. Foley catheter versus vaginal prostaglandin E2 gel for induction of labour at term (PROBAAT trial): an open-label, randomised controlled trial. *Lancet* 2011; 378: 2095–103.
- 101 Alfrevic Z, Keeney E, Dowswell T, et al. Labour induction with prostaglandins: a systematic review and network meta-analysis. *BMJ* 2015; 350: h217.
- 102 Broekhuijsen K, van Baaren GJ, Van Pampus MG, et al, and the HYPITAT-II study group. Immediate delivery versus expectant monitoring for hypertensive disorders of pregnancy between 34 and 37 weeks of gestation (HYPITAT-II): a multicentre, open-label, randomised controlled trial. *Lancet* 2015; 385: 2492–501.
- 103 Vigil-De Gracia P, Reyes Tejada O, Calle Miñaca A, et al. Expectant management of severe preeclampsia remote from term: the MEXPRE Latin Study, a randomized, multicenter clinical trial. *Am J Obstet Gynecol* 2013; 209: 425. e1–8.
- 104 Rana S, Powe CE, Salahuddin S, et al. Angiogenic factors and the risk of adverse outcomes in women with suspected preeclampsia. *Circulation* 2012; 125: 911–19.
- 105 Brownfoot FC, Gagliardi DI, Bain E, Middleton P, Crowther CA. Different corticosteroids and regimens for accelerating fetal lung maturation for women at risk of preterm birth. *Cochrane Database Syst Rev* 2013; 8: CD006764.
- 106 Gyamfi-Bannerman C, Gilbert S, Landon MB, et al, and the Eunice Kennedy Shriver National Institute of Child Health, and the Human Development (NICHD) Maternal-Fetal Medicine Units Network (MFMU). Effect of antenatal corticosteroids on respiratory morbidity in singletons after late-preterm birth. *Obstet Gynecol* 2012; 119: 555–59.

- 107 Doyle LW, Crowther CA, Middleton P, Marret S, Rouse D. Magnesium sulphate for women at risk of preterm birth for neuroprotection of the fetus. *Cochrane Database Syst Rev* 2009; 1: CD004661.
- 108 Epstein FH. Late Vascular Effects of Toxemia of Pregnancy. *N Engl J Med* 1964; 271: 391–95.
- 109 Cusimano MC, Pudwell J, Roddy M, Cho CK, Smith GN. The maternal health clinic: an initiative for cardiovascular risk identification in women with pregnancy-related complications. *Am J Obstet Gynecol* 2014; 210: 438. e1–9.
- 110 Sattar N, Bedomir A, Berry C, Shepherd J, Greer IA, Packard CJ. Lipoprotein subfraction concentrations in preeclampsia: pathogenic parallels to atherosclerosis. *Obstet Gynecol* 1997; 89: 403–08.
- 111 Kvehaugen AS, Dechend R, Ramstad HB, Troisi R, Fugelseth D, Staff AC. Endothelial function and circulating biomarkers are disturbed in women and children after preeclampsia. *Hypertension* 2011; 58: 63–69.
- 112 McDonald SD, Malinowski A, Zhou Q, Yusuf S, Devereaux PJ. Cardiovascular sequelae of preeclampsia/eclampsia: a systematic review and meta-analysis. *Am Heart J* 2008; 156: 918–30.
- 113 Hermes W, Franx A, van Pampus MG, et al. Cardiovascular risk factors in women who had hypertensive disorders late in pregnancy: a cohort study. *Am J Obstet Gynecol* 2013; 208: 474. e1–8.
- 114 Veerbeek JH, Hermes W, Breimer AY, et al. Cardiovascular disease risk factors after early-onset preeclampsia, late-onset preeclampsia, and pregnancy-induced hypertension. *Hypertension* 2015; 65: 600–06.
- 115 Williams D. Pregnancy: a stress test for life. *Curr Opin Obstet Gynecol* 2003; 15: 465–71.
- 116 Jacobs AK, Eckel RH. Evaluating and managing cardiovascular disease in women: understanding a woman's heart. *Circulation* 2005; 111: 383–84.
- 117 McDonald SD, Han Z, Walsh MW, Gerstein HC, Devereaux PJ. Kidney disease after preeclampsia: a systematic review and meta-analysis. *Am J Kidney Dis* 2010; 55: 1026–39.
- 118 Hoedjes M, Berks D, Vogel I, et al. Effect of postpartum lifestyle interventions on weight loss, smoking cessation, and prevention of smoking relapse: a systematic review. *Obstet Gynecol Surv* 2010; 65: 631–52.
- 119 Heidrich MB, Wenzel D, von Kaisenberg CS, Schippert C, von Versen-Höyneck FM. Preeclampsia and long-term risk of cardiovascular disease: what do obstetrician-gynecologists know? *BMC Pregnancy Childbirth* 2013; 13: 61.
- 120 MacDonald SE, Walker M, Ramshaw H, Godwin M, Chen XK, Smith G. Hypertensive disorders of pregnancy and long-term risk of hypertension: what do Ontario prenatal care providers know, and what do they communicate? *J Obstet Gynaecol Can* 2007; 29: 705–10.
- 121 Young B, Hacker MR, Rana S. Physicians' knowledge of future vascular disease in women with preeclampsia. *Hypertens Pregnancy* 2012; 31: 50–58.
- 122 Cusimano MC, Pudwell J, Roddy M, et al. The Maternal Health Clinic: an initiative for cardiovascular risk identification in women with pregnancy-related complications. *Am J Obstet Gynecol* 2014; 210: 438. e1–9.
- 123 Spratling PM, Pryor ER, Moneyham LD, Hodges AL, White-Williams CL, Martin JN Jr. Effect of an educational intervention on cardiovascular disease risk perception among women with preeclampsia. *J Obstet Gynecol Neonatal Nurs* 2014; 43: 179–89.
- 124 Berks D, Hoedjes M, Raat H, Duvekot JJ, Steegers EA, Habbema JD. Risk of cardiovascular disease after pre-eclampsia and the effect of lifestyle interventions: a literature-based study. *BJOG* 2013; 120: 924–31.
- 125 Rautio N, Jokelainen J, Korpi-Hyövälti E, et al. Lifestyle intervention in prevention of type 2 diabetes in women with a history of gestational diabetes mellitus: one-year results of the FIN-D2D project. *J Womens Health (Larchmt)* 2014; 23: 506–12.
- 126 Ahmed R, Dunford J, Mehran R, Robson S, Kunadian V. Pre-eclampsia and future cardiovascular risk among women: a review. *J Am Coll Cardiol* 2014; 63: 1815–22.
- 127 Firoz T, Sanghvi H, Merialdi M, von Dadelszen P. Pre-eclampsia in low and middle income countries. *Best Pract Res Clin Obstet Gynaecol* 2011; 25: 537–48.
- 128 Dunsmuir DT, Payne BA, Cloete G, et al. Development of mHealth applications for pre-eclampsia triage. *IEEE J Biomed Health Inform* 2014; 18: 1857–64.
- 129 Villar J, Ba'aqeel H, Piaggio G, et al, and the WHO Antenatal Care Trial Research Group. WHO antenatal care randomised trial for the evaluation of a new model of routine antenatal care. *Lancet* 2001; 357: 1551–64.
- 130 Vogel JP, Habib NA, Souza JP, et al. Antenatal care packages with reduced visits and perinatal mortality: a secondary analysis of the WHO Antenatal Care Trial. *Reprod Health* 2013; 10: 19.
- 131 Hofmeyr GJ, Hodnett ED. Antenatal care packages with reduced visits and perinatal mortality: a secondary analysis of the WHO antenatal care trial - Commentary: routine antenatal visits for healthy pregnant women do make a difference. *Reprod Health* 2013; 10: 20.
- 132 Cormick G, Zhang NN, Andrade SP, et al. Gaps between calcium recommendations to prevent pre-eclampsia and current intakes in one hospital in Argentina. *BMC Res Notes* 2014; 7: 920.
- 133 Belizán JM, Villar J. The relationship between calcium intake and edema-, proteinuria-, and hypertension-getosis: an hypothesis. *Am J Clin Nutr* 1980; 33: 2202–10.
- 134 Hofmeyr GJ, Belizán JM, von Dadelszen P, and the Calcium and Pre-eclampsia (CAP) Study Group. Low-dose calcium supplementation for preventing pre-eclampsia: a systematic review and commentary. *BJOG* 2014; 121: 951–57.
- 135 Goldenberg RL, McClure EM, MacGuire ER, Kamath BD, Jobe AH. Lessons for low-income regions following the reduction in hypertension-related maternal mortality in high-income countries. *Int J Gynaecol Obstet* 2011; 113: 91–95.
- 136 Althabe F, Belizán JM, McClure EM, et al. A population-based, multifaceted strategy to implement antenatal corticosteroid treatment versus standard care for the reduction of neonatal mortality due to preterm birth in low-income and middle-income countries: the ACT cluster-randomised trial. *Lancet* 2015; 385: 629–39.
- 137 Goldenberg RL, Jones B, Griffin JB, et al. Reducing maternal mortality from preeclampsia and eclampsia in low-resource countries—what should work? *Acta Obstet Gynecol Scand* 2015; 94: 148–55.
- 138 Steegers EA, von Dadelszen P, Duvekot JJ, Pijnenborg R. Pre-eclampsia. *Lancet* 2010; 376: 631–44.
- 139 Comet Initiative. Developing a core outcome set for hypertensive disorders in pregnancy. <http://www.comet-initiative.org/studies/details/588> (accessed April 7, 2015).
- 140 Myatt L, Redman CW, Staff AC, et al, and the Global Pregnancy CoLaboratory. Strategy for standardization of preeclampsia research study design. *Hypertension* 2014; 63: 1293–301.
- 141 Burton GJ, Sebire NJ, Myatt L, et al. Optimising sample collection for placental research. *Placenta* 2014; 35: 9–22.
- 142 Mol BW, Ruifrok AE, and the Global Obstetrics Network. Global alignment, coordination and collaboration in perinatal research: the Global Obstetrics Network (GONet) Initiative. *Am J Perinatol* 2013; 30: 163–66.